



Severity and impact of Douglas-fir beetle infestations in northern Wyoming

Kurt K. Allen, Denise M. Hardesty and Daniel F. Long, USDA Forest Service R2 Forest Health Management, 1730 Samco Road, Rapid City, SD 57702, USA



INTRODUCTION

The Douglas-fir beetle (DFB) *Dendroctonus pseudotsugae* infests and kills Douglas-fir throughout its range in the west. Typically, the Douglas-fir beetle reproduces in scattered Douglas-fir (*Pseudotsuga menziesii*) that are highly stressed, such as windfall, defoliated or fire-scorched trees (Furniss 1962, Furniss 1965). If enough suitable host material is present, the beetles can infest nearby healthy trees. Previous research on Douglas-fir beetle has looked at forest stand conditions and site characteristics that lead to beetle infestations (Furniss 1965, Furniss et al. 1979, Negron 1998, Negron et al. 1999). Also, we have previously looked at the impacts on forest overstory and understory conditions in areas attacked by Douglas-fir beetle 4-5 years prior to study (McMillin and Allen 2003). Significant changes to both the overstory and understory conditions were documented.

Aerial and ground surveys indicate a rapidly increasing Douglas-fir beetle outbreak in the Bighorn Mountains of northern Wyoming (Figure 1). While the beetle typically reproduces in stressed trees, no stressor such as blowdown or fire is evident to have initiated this outbreak.

The objectives of this study are to: 1. Quantify the severity and impacts of Douglas-fir beetle infestations at the stand level through a plot network on the Bighorn National Forest. This includes both overstory and understory changes. 2. Describe in detail the forest conditions present in areas that have experienced high levels of mortality over the past few years. 3. Provide information to forest managers and planners in the northern Rocky Mountains on potential impacts of Douglas-fir beetle based on stand conditions examined.



Figure 1. Douglas-fir tree mortality caused by Douglas-fir beetle on the Bighorn National Forest, WY in the (A)Tensleep Canyon and (B)Shell Canyon areas.

METHODS

56 plots were installed in stands that are predominantly Douglas-fir throughout three areas along the western edge of the Bighorn National Forest in north central Wyoming: Shell Canyon, Battle Park, and Tensleep Canyon (Figure 2). Fixed radius plots were installed in stands that had been impacted by Douglas-fir beetle and those that were currently unaffected. Within each plot the following measurements were taken on all trees >4 inches diameter at breast height (DBH): tree species, DBH, crown class and condition/damages. Year of attack was determined for those trees colonized by Douglas-fir beetle. The species and number of established seedlings (6 inches and taller) and saplings were counted within an 11.8 foot radius plot (1/100th acre) around plot center. Line transects were used to determine understory conditions. The number of total inches covered by grass, forb and shrub species was measured along a 10 foot line north, south, east and west of plot center. Understory canopy height was measured at its highest point along each of the cardinal directions.




Figure 2. Location of study sites on the Bighorn National Forest WY.

RESULTS

Figure 3 shows the results of aerial surveys done in 2001 and 2004 and the subsequent change in tree mortality caused by Douglas-fir beetle. The outbreak has increased dramatically in this 3 year period. Table 1 lists the preliminary results from the infested and uninfested plots measured in the summer of 2004. Although the results have not been statistically analyzed yet, there are some obvious differences in stand conditions before and after Douglas-fir beetle infestations and between infested and uninfested plots. DBH, % Douglas-fir, and Douglas-fir basal area all decreased in infested plots. Figure 4 shows this is a relatively new outbreak since most trees have been killed in the last 1-3 years. There are some minor changes in understory conditions (Figure 5) although these may become more pronounced the longer the infestation continues.




Figure 3. Comparison of the extent of Douglas-fir beetle activity in the Bighorn National Forest, WY between 2001 and 2004 based on aerial detection surveys.

Table 1. Summary (mean +/- SEM) of Douglas-fir beetle impacts on stand conditions

Variable	Uninfested	Infested
Total Basal Area (ft ² /ac)	221 (13.9)	298 (16.0)
Live Basal Area (ft ² /ac)	216 (13.3)	66 (45.6)
Total DBH (inches)	10.9 (.4)	13.8 (.7)
Live DBH (inches)	10.9 (.4)	8.0 (.7)
Percent DF (pre-infested)	85.1 (3.1%)	85.8 (3.3)
Percent DF (post-infested)	84.3 (2.9%)	57.0 (7.4)
Understory Height (inches)	7.6 (1.1)	9.7 (.9)
Regeneration (per 0.01 ac plot)	2.14 (.4)	2.39 (.6)

CONCLUSIONS

1. There has been and continues to be a rapidly increasing Douglas-fir beetle outbreak occurring in these areas.
2. This outbreak has the potential to significantly change forest conditions in the Douglas-fir type.
3. The minor changes already showing in the understory are likely to increase as the recently killed trees lose needles and new mortality in the overstory occurs.

PLANS

- Revisit the plots in August of 2005 to measure new Douglas-fir beetle caused mortality.
- Analyze remotely sensed data using an unsupervised classification process to study the extent and spread of Douglas-fir beetle epidemic in the study areas.
- Compile and statistically analyze the data.
- Prepare reports for managers' use on impacts of Douglas-fir beetle outbreaks and the potential changes in forest vegetation.




Figure 4. Tree mortality caused by Douglas-fir beetle on the Bighorn National Forest per year by basal area and number of trees. Live trees=Surviving Douglas-fir as of 2004.




Figure 5. Impact of Douglas-fir beetle on understory ground cover in the Bighorn National Forest, WY.

Literature Cited

- Furniss, M.M., 1962. Infestation patterns of Douglas-fir beetle in standing and windthrown trees in southern Idaho. *J. Econ. Entomol.*, 55:486-491.
- Furniss, M.M., 1965. Susceptibility of fire-injured Douglas-fir to bark beetle attack in southern Idaho. *J. For.*, 63: 8-11.
- Furniss, M.M., McGregor, M.D., Foiles, M.W. and Partridge, A.D., 1979. Chronology and characteristics of a Douglas-fir beetle outbreak in northern Idaho. USDA For. Serv. Gen. Tech. Rep. INT-59, 19 pp.
- McMillin, J.D. and K.K. Allen, 2003. Effects of Douglas-fir beetle (Coleoptera: Scolytidae) infestations on forest overstory and understory conditions in western Wyoming. *West. N. Amer. Natur.*, 63: 498-506.
- Negron, J.F., 1998. Probability of infestation and extent of mortality associated with the Douglas-fir beetle in the Colorado Front Range. *For. Ecol. Manage.*, 107: 71-85.
- Negron, J.F., Schupp, Jr., W.C., Gibson, K.E., Anhold, J., Hansen, D., Thier, R. and Mocellini, P., 1999. Estimating extent of mortality associated with the Douglas-fir beetle in the central and northern Rockies. *West. J. Appl. For.*, 14: 121-127.